



# Association of Gastritis with Sleep and Quality of Life: A Hospital-based Cross-sectional Study

## Gastritin Uyku ve Yaşam Kalitesi ile İlişkisi: Hastane Temelli Kesitsel Bir Çalışma

Hasan Cantay, Ece Büyüksandalyacı Tunç\*

Kafkas University Faculty of Medicine, Department of General Surgery, Kars, Turkey

\*Kars Harakani State Hospital, Clinic of Psychiatry, Kars, Turkey

### Abstract

**Objective:** Gastritis is an important public health problem that negatively affects many people in the world. In studies conducted, the most frequently mentioned risk factors for gastritis in studies are *H. pylori*, spicy and smoked foods, smoking, and alcohol consumption. This study investigates the effect of life and sleep quality on gastritis, which has not been emphasized much before.

**Materials and Methods:** In this hospital-based cross-sectional study, 277 participants who underwent gastroscopy were given the Pittsburgh sleep quality scale and the quality of life index. Statistical analysis of the data was done with SPSS 22.0.

**Results:** Gastritis was detected in 66.4% of the patients who underwent gastroscopy. According to the logistic regression analysis results, gastritis is 6.935 (confidence interval: 3.023-15.908) times more common in people with poor sleep quality than in people with good sleep quality. We determined that the probability of developing gastritis was high in those with poor sleep quality and mental health.

**Conclusion:** It is important to identify risk factors for the prevention and early treatment of gastritis. According to the findings of this study, it is thought that stomach complaints should be specifically questioned in patients with sleep and mental disorders.

**Keywords:** Gastritis, mental health, quality of life, sleep disorders, stress

### Öz

**Amaç:** Gastrit, dünyada birçok insanı olumsuz etkileyen önemli bir halk sağlığı sorunudur. Yapılan çalışmalarda gastrit için en sık belirtilen risk faktörlerinin *H. pylori*, baharatlı ve tütülenmiş yiyecekler, sigara ve alkol tüketimi olduğu gösterilmiştir. Bu çalışma, yaşam ve uyku kalitesinin gastrit üzerine daha önce üzerinde fazla durulmayan etkisini araştırmayı amaçlamaktadır.

**Gereç ve Yöntem:** Hastane temelli bu kesitsel çalışmada gastroskopi yapılan 277 katılımcıya Pittsburgh uyku kalitesi ölçeği ve yaşam kalitesi indeksi verildi. Verilerin istatistiksel analizi SPSS 22.0 ile yapıldı.

**Bulgular:** Gastroskopi yapılan hastaların %66,4'ünde gastrit saptandı. Lojistik regresyon analizi sonuçlarına göre uyku kalitesi kötü olan kişilerde gastrit, uyku kalitesi iyi olan kişilere göre 6,935 (güven aralığı: 3,023-15,908) kat daha fazla görülmekte idi. Uyku kalitesi ve ruh sağlığı kötü olanlarda gastrit gelişme olasılığının yüksek olduğu belirlendi.

**Sonuç:** Gastritin önlenmesi ve erken tedavisi için risk faktörlerinin belirlenmesi önemlidir. Bu çalışmada, gastrite neden olabilecek uyku ve ruhsal bozuklukları olan hastalarda mide şikayetlerinin özel olarak sorgulanması gerektiği sonucuna varılmıştır.

**Anahtar Kelimeler:** Gastrit, ruh sağlığı, yaşam kalitesi, uyku bozuklukları, stres

### Introduction

Gastritis is a widespread disease that can last a lifetime, progresses confidentially in humans, and indicates inflammation of the gastric mucosa (1). It is the most common gastroscopy finding in patients presenting with dyspepsia, nausea, and vomiting (2). Gastritis is still a serious public health problem in both developed and developing countries. It is a disease that affects individuals' socio-economic status, lifestyles, living conditions, behaviors, and habits, in short, their living standards. 50.8% of the population in developing countries and 34.7% of the population in developed countries have health problems

due to gastritis. In this context, gastritis threatens patients' physical health and impairs their quality of life (QoL) (3).

Previous studies have revealed that factors such as gender, age, socio-economic status, biological, environmental factors, and individual behaviors may contribute to gastritis. The most common cause of gastritis is *Helicobacter pylori* infection (4,5). Other causes include eating acidic, spicy, or hot meals often, drinking too much alcohol, smoking, and taking certain medications like nonsteroidal anti-inflammatory drugs (NSAIDs). Chronic stress and impairment of mental health can also induce gastritis (6-8). Although gastric acid is essential for the digestion

Address for Correspondence/Yazışma Adresi: Hasan Cantay MD, Kafkas University Faculty of Medicine, Department of General Surgery, Kars, Turkey

Phone: +90 533 623 55 76 E-mail: hasan\_cantay@hotmail.com ORCID-ID: orcid.org/0000-0003-3309-8879

Received/Geliş Tarihi: 09.12.2021 Accepted/Kabul Tarihi: 20.05.2022

©Copyright 2022 by Turkish Sleep Medicine Society / Journal of Turkish Sleep Medicine published by Galenos Publishing House.

of food, in cases of psychological stress, gastric acid secretion increases and prepares the ground for gastritis (7,9). Gastritis has been reported many times in the literature to impair sleep and life quality (5,10,11). However, when we look at the matter from the opposite side, a few studies have shown that poor sleep quality, causes an increase in gastrointestinal diseases. Excessive secretion of proinflammatory cytokines in sleep disorders causes gastric mucosa to be damaged more easily (12,13). Dyspepsia, which is one of the most important symptoms of gastritis, can be seen due to sleep disorders and related changes in eating habits (14).

There are few studies on psychiatric conditions that may cause gastritis such as depressive disorder and anxiety disorder (8,15,16). As far as we know, sleep disorders and poor mental health affecting the QoL that can cause gastritis have not been studied sufficiently. Aware of such gaps in the work environment and limited evidence, this study aimed to evaluate the implications of poor mental health and sleep quality on gastritis. Identifying the risk factors specifically associated with gastritis is an essential point to implement the necessary measures. It is therefore vital to distinguish the factors that may cause gastritis and, implement measures to intervene against factors that contribute to gastritis.

## Materials and Methods

**Research design:** This is a hospital-focused cross-sectional study.

**The research universe:** The number of gastroscopies performed in the general surgery endoscopy unit in 2020 as the universe in the research was formed. The number of gastroscopies performed in 2020 was 998. It is assumed that the same number of gastroscopies will be performed in 2021.

**The sample of the study:** The number of people to be sampled was calculated with the formula  $n = Nt^2 p q / d^2 (N-1) + t^2 p q$ . N is the number of individuals in the universe; n is the number of individuals to be sampled; p is the incidence (probability) of the event under investigation; q is the frequency (probability) of the event under investigation; t is the theoretical value found in the table t at a given degree of freedom and the detected error level; d is the desired  $\pm$  deviation according to the incidence of the event. Accordingly,  $p=0.50$ ;  $q=0.50$ ;  $t=1.96$ ; when  $d=0.05$ , the sample size was determined as 277 people.

**Inclusion criteria for the study:** Patients who were diagnosed with gastritis as a result of the gastroscopy procedure and accepted to participate in the study.

**Exclusion criteria from the study:** Cases with normal gastroscopic findings or malignancy as a result of the gastroscopy procedure.

**Verbal/written consent and ethics committee approval:** After the Kafkas University Faculty of Medicine Ethics Committee approval (date: 11.03.2020 and no: 80576354-050-99/19), written and verbal consent was obtained from the patients. Afterward, the data of the study were collected.

**Structuring the data collection form:** The data collection form of the study was prepared by the researchers.

**The dependent variable of the study:** Gastritis status in people undergoing gastroscopy in the general surgery endoscopy unit. The study's independent variables are age, gender, Pittsburgh sleep quality index (PSQI), and QoL index.

**The preliminary trial of the study:** It was conducted with five patients who applied to the outpatient clinic and would not be included in the study. After the preliminary test, necessary corrections were made in the data collection form.

**Data collection:** Data was collected between April-October 2021 by the general surgeon and psychiatrist who conducted the research using the face-to-face interview technique.

## Measurements

**The PSQI:** Comprises 19 items and a widely used self-reported questionnaire. It is regarded as a general tool for measuring sleep quality. Its subcomponents are; subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, usage of sleep medication, and daytime dysfunction. A score of 5 or above indicates bad sleep quality (17). The Turkish validity and reliability study of the Pittsburg sleep quality was done by Agargun et al. (18).

**Short-form health survey (SF-36):** SF-36 was prepared by Rand Corporation in 1992 as short form-36 (SF-36). It, provides a wide-angle measurement among QoL scales. The scale includes items that include the perception of change in health in the last four weeks and the perception of change in health in the last week. SF-36 is a short, suitable for use in clinical practice and research. However, it has a strong general overview in terms of its comprehensive, psychometric properties. As the name of the scale suggests, 36 items consist of physical and mental components. It has three main topics (functional status, well-being, general understanding of health) and 8 health concepts (physical function, social function, physical role function, emotional role function, mental health, vitality, fatigue, pain, general perspective) (19). SF-36 was first adapted to Turkish society by Kocyigit (20), after validation.

## Statistical Analysis

Statistical analysis of the study was done with SPSS 22.0 computer program. Student t-test and chi-square tests were used in pairwise analyses. Statistically significant variables ( $p<0.05$ ) were included in the logistic regression analysis.

## Results

Two hundred seventy-seven people participated in the study. One hundred thirty-nine (50.2%) of the participating individuals were under 47 years old, 138 (49.8%) were 48 years old and over. One hundred sixty (57.8%) were female and 117 (43.2%) were male. 66.4% of the people who underwent gastroscopy were diagnosed with gastritis.

When we look at the relationship between sleep quality and gastritis in Table 1, gastritis was found to be significantly more common in those with poor subjective sleep quality, sleep latency, habitual sleep efficiency, sleep disturbances, and daytime dysfunction ( $p<0.05$ ). No significant difference was found in terms of sleep duration and usage of sleep medication in patients with gastritis ( $p>0.05$ ).

According to the relationship between QoL and gastritis in Table 2, no significant relationship was found between physical and social function subscales and gastritis ( $p>0.05$ ). A highly significant correlation was found between those with low total scores and all subscale scores except physical and social subscales, and gastritis ( $p<0.05$ ).

According to the logistic regression analysis results, gastritis is 6.935 [confidence interval (CI): 3.023-15.908] times more common in people with poor sleep quality than people with good sleep quality. On the other hand, while the effect of physical health on QoL is lost in the regression analysis, gastritis is 1.163 (CI: 1.112-1.216) times more common in those with

poor mental health than in those with good mental health (Table 3).

## Discussion

When we look at the studies conducted both in the world and in Turkey, although many data have been obtained about the causes and consequences of gastritis, the number of studies on the effects of poor mental health and sleep quality on gastritis, which significantly reduces the QoL, is limited. Our study aims to contribute to the gap in the literature that impaired sleep quality and mental health may cause gastritis. In this way, it is important to question the mental health and sleep quality of

Subcomponents	Gastritis	n (%)	Mean	Standard deviation ( $\pm$ )	p
Subjective sleep quality	Yes	184 (66.4)	1.02	0.71	0.001
	No	93 (33.6)	0.46	0.50	
Sleep latency	Yes	184 (66.4)	1.08	0.81	0.001
	No	93 (33.6)	0.26	0.51	
Sleep duration	Yes	184 (66.4)	0.58	0.79	0.096
	No	93 (33.6)	0.40	0.92	
Habitual sleep efficiency	Yes	184 (66.4)	0.61	0.80	0.029
	No	93 (33.6)	0.39	0.79	
Sleep disturbances	Yes	184 (66.4)	1.35	0.52	0.001
	No	93 (33.6)	1.02	0.33	
Usage of sleep medication	Yes	184 (66.4)	0.56	0.74	0.116
	No	93 (33.6)	0.41	0.77	
Daytime dysfunction	Yes	184 (66.4)	3.01	10.60	0.044
	No	93 (33.6)	0.75	0.67	
Total score	Yes	184 (66.4)	6.91	2.96	0.001
	No	93 (33.6)	3.69	1.75	

Subcomponents	Gastritis	n (%)	Mean	Standard deviation ( $\pm$ )	p
Physical function	Yes	184 (66.4)	63.70	22.38	0.118
	No	93 (33.6)	68.01	19.99	
Difficulty with physical role	Yes	184 (66.4)	41.28	25.64	0.001
	No	93 (33.6)	76.88	20.93	
Pain	Yes	184 (66.4)	46.22	16.69	0.001
	No	93 (33.6)	67.85	10.62	
General health	Yes	184 (66.4)	43.32	9.59	0.001
	No	93 (33.6)	57.74	10.18	
Vitality	Yes	184 (66.4)	46.44	8.87	0.001
	No	93 (33.6)	62.10	11.40	
Social function	Yes	184 (66.4)	61.62	21.03	0.292
	No	93 (33.6)	64.11	12.26	
Emotional role	Yes	184 (66.4)	29.89	25.67	0.001
	No	93 (33.6)	86.74	17.82	
Mental health	Yes	184 (66.4)	42.46	11.44	0.001
	No	93 (33.6)	61.51	9.23	

**Table 3. The effect of age, gender, sleep quality, physical and mental health on gastritis and logistic regression results**

Independent variables		Gastritis	No gastritis	Gastritis	No gastritis	p	OR	95% CI (EK-EB value)*	p
		N (%)	N (%)	Mean (± SD)*	Mean (± SD)*				
Age	≤47	85 (61.2)	54 (38.8)			0.062			
	≥48	99 (71.7)	39 (28.3)						
Gender	Woman	111 (69.4)	49 (30.6)			0.224			
	Man	73 (62.4)	44 (37.6)						
Sleep quality	Good	18 (20.7)	69 (79.3)			0.001	0.144	0.063-0.331	0.001
	Bad	166 (87.4)	24 (12.6)						
Physical health				48.63±14.19	67.62±9.11	0.001	0.971	0.931-1.012	0.166
Mental health				45.10±12.24	68.61±8.48	0.001	0.860	0.822-0.899	0.001

CI: Confidence interval, SD: Standard deviation, OR: Odds ratio

patients diagnosed with gastritis, and also, to emphasize the importance of questioning gastric complaints in those with sleep disorders and poor mental health.

In this study, patients with nausea, vomiting, stomach pain, and heartburn complaints underwent a gastroscopy and 66.4% of these patients were diagnosed with gastritis. Gastroscopy is the most commonly used study for the diagnosis of gastric pathologies (4). Gastritis is an important public health problem. 50.8% of the population in developing countries suffer from gastritis. 54.8% of adults who applied to health institutions in Kenya were clinically diagnosed with gastritis. In Ethiopia, reported that 53% of individuals between the ages of 54-61 had gastritis due to *H. pylori* infection. The result showed that the prevalence of gastritis was 73.5% in a study conducted in Pakistan (21,22). Since Turkey is a developing country, our results are consistent with the literature. It can be thought that our study can represent the whole country.

In this study, gastritis was found 6.935 (CI: 3.023-15.908) times more in those with low sleep quality than in those with high sleep quality. According to literature, sleep disorders may cause gastrointestinal system diseases such as gastroesophageal reflux, peptic ulcer, irritable bowel disease, inflammatory bowel disease, gastric and colorectal cancers. However, studies on gastritis are insufficient (23-26). Sleep quality affects the pathogenesis and emergence of digestive system diseases. The healthy circadian rhythm and sleep functions in the gastrointestinal system are intestinal stem cell proliferation, motility regulation, protein and carbohydrates digestion and absorption, electrolyte balance, intestinal microbiota protection, and intestinal barrier protection. Sleep is regulated by the light-dark cycle and, accordingly, by the circadian rhythm. Individually differential regulation or variation of the circadian rhythm is a common cause of sleep complaints, which plays an important role in the occurrence of many gastrointestinal disorders (25,26). In sleep disorders, pro-inflammatory cytokines that make the gastric mucosa unprotected and fragile increase, and lead to gastritis (12,13). The eating habits of individuals with sleep disorders also change. Therefore, the incidence of gastritis increases in these individuals (14). Although studies on this subject are few, it is reported that gastritis is associated with low sleep quality

(27,28). Gastritis is more common in studies conducted with individuals who work night shifts and have disturbed sleep patterns (29,30). According to these findings and our results, it can be said that low sleep quality may cause gastritis.

In this study, gastritis was found 1.163 (CI: 1.112-1.216) times more common in those with poor mental health than in those with good mental health. The situation that can be confusing here is the question of whether gastritis may cause mental problems. In the QoL index we used in our study, the patients' mental problems for the last 1 month are questioned. In psychiatric examinations, it was observed that these patients struggled with mental problems for more than 1 month, and gastric complaints appeared later. In this context, poor mental health was thought to cause gastritis. Stress which indicates a deterioration in mental health causes excessive secretion of gastric acid, which is normally required for the digestion of food, and leads to deterioration of integrity of the stomach wall and causes gastritis (31). Also, for the gastric mucosa to be healthy, the production of hydrogen sulfide (H<sub>2</sub>S) must be sufficient and the gastric flora must be intact. Vitamin B6 plays a leading role in the production of H<sub>2</sub>S in the stomach. In individuals under psychological stress, vitamin B6 level decreases, H<sub>2</sub>S production in the stomach decreases, and the gastric flora deteriorate. In this case, the integrity of the gastric mucosa is disturbed and gastritis is observed (7,32). According to studies, high-stress levels, anxiety and depression are associated with gastritis (8,27,33-35). Another study reported that the probability of gastritis is high in those with poor mental health (36). In a study conducted with young people, it was found that stress increases especially during seasonal exam periods, which increases the incidence of gastritis (37). Although these results should be supported by more studies on this subject, it can be said that the risk of gastritis increases in individuals with impaired mental health, both by the information in the literature and our findings. Since gastritis affects a significant part of the population, identifying the risk factors that cause gastritis can prevent gastritis. It is important to identify risk factors based on patients. Although studies conducted so far have found various risk factors, studies investigating the effects of mental health and sleep quality on gastritis are limited. According to our study,



gastritis is more common in people with impaired sleep quality and mental health. Therefore, it is very important to question gastric complaints in patients with these problems, both to prevent gastritis and to provide early diagnosis and treatment. Along with these results of our study, it should be kept in mind that gastritis may also affect sleep and QoL.

### Study Limitations

One of the limitations of our study can be considered that the number of cases is relatively low, therefore more multicenter case series in future studies will lead to more inclusive results. Another limitation is that the eating habits and body mass index of the patients are not taken into account. It is known that if the BMI is high, sleep quality decreases considerably, and gastritis is more common in these cases (38). Therefore, it can be done by considering BMI and eating habits in future studies. Another limitation is that the physical exercise status of the individuals was not taken into account. Regular physical exercise is protective in terms of mental health, sleep patterns, and gastrointestinal diseases. Therefore, physical exercise status should be considered in future studies.

### Conclusion

Since it is known that drugs such as NSAIDs can cause gastrointestinal diseases, medications used by the patients should also be questioned in future studies. In our study, patients were evaluated with the PSQI and SF-36 tests. The inability to evaluate patients with anxiety and depression scales is another important limitation of the study.

### Acknowledgments

We thank Associate Professor Binali Çatak for helping the statistical analyses.

### Ethics

**Ethics Committee Approval:** After the Kafkas University Faculty of Medicine Ethics Committee approval (date: 11.03.2020 and no: 80576354-050-99/19).

**Informed Consent:** Written and verbal consent was obtained from the patients.

**Peer-review:** Internally and externally peer-reviewed.

### Authorship Contributions

Concept: H.C., E.B.T., Data Collection or Processing: H.C., E.B.T., Analysis or Interpretation: H.C., Funding: H.C., Writing: H.C., E.B.T.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

### References

1. Megha R, Farooq U, Lopez PP. Stress-Induced Gastritis. 2021 Aug 31. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021. PMID: 29763101
2. Marcial G, Rodríguez C, Medici M, Valdez GF. New Approaches in Gastritis Treatment. *Gastritis and Gastric Cancer - New Insights in Gastroprotection, Diagnosis and Treatments* 2012;153-76.
3. Wen Z, Li X, Lu Q, Brunson J, Zhao M, Tan J, Wan C, Lei P. Health related quality of life in patients with chronic gastritis and peptic ulcer and factors with impact: A longitudinal study. *BMC Gastroenterol* 2014;14:149.
4. Rugge M, Savarino E, Sbaraglia M, Bricca L, Malfertheiner P. Gastritis: The clinico-pathological spectrum. *Dig Liver Dis* 2021;53:1237-46.
5. Takeoka A, Tayama J, Kobayashi M, Sagara I, Ogawa S, Saigo T, Hayashida M, Yamasaki H, Fukudo S, Shirabe S. Psychological effects of *Helicobacter pylori*-associated atrophic gastritis in patients under 50 years: A cross-sectional study. *Helicobacter* 2017;22. doi: 10.1111/HEL.12445
6. Lee SP, Sung IK, Kim JH, Lee SY, Park HS, Shim CS. The effect of emotional stress and depression on the prevalence of digestive diseases. *J Neurogastroenterol Motil* 2015;21:273-82.
7. Han Y, Li Y, Hu Z, Wang X, Ren X, Yu Y, Li Y, Li W, Sun Y. Chronic psychological stress induces gastritis in C57BL/6 mice with decreased production of hydrogen sulphide (H2S) and microbial dysbiosis. (2019). Available at SSRN 3360097.
8. Wau ET, Amidos J, Simamora PM. Levels of Stress Related to Incidence of Gastritis in Adolescents. *Mental Health* 2018;4:2.
9. Auguste LJ, Lackner R, Ratner L, Stein TA, Bailey B. Prevention of stress-induced erosive gastritis by parenteral administration of arachidonic acid. *JPEN J Parenter Enteral Nutr* 1990;14:615-7.
10. Magni G, Salmi A, Paterlini A, Merlo A. Psychological distress in duodenal ulcer and acute gastroduodenitis. A controlled study. *Dig Dis Sci* 1982;27:1081-4.
11. Solov'eva AD, Sheptulin AA, Annamamedova R. [Emotional-personality condition of patients with some gastroenterological diseases]. *Klin Med (Mosk)* 1997;75:27-8.
12. Ali T, Choe J, Awab A, Wagener TL, Orr WC. Sleep, immunity and inflammation in gastrointestinal disorders. *World J Gastroenterol* 2013;19. doi: 10.3748/wjg.v19.i48.9231.
13. Khanijow V, Prakash P, Emsellem HA, Borum ML, Doman DB. Sleep Dysfunction and Gastrointestinal Diseases. *Gastroenterol Hepatol (N Y)* 2015;11:817-25.
14. Miwa H. Why dyspepsia can occur without organic disease: Pathogenesis and management of functional dyspepsia. *J Gastroenterol* 2012;47:862-71.
15. Goodwin RD, Cowles RA, Galea S, Jacobi F. Gastritis and mental disorders. *J Psychiatr Res* 2013;47:128-32.
16. Painsipp E, Wultsch T, Shahbazian A, Edelsbrunner M, Kreissl MC, Schirbel A, Bock E, Pabst MA, Thoeringer CK, Huber HP, Holzer P. Experimental gastritis in mice enhances anxiety in a gender-related manner. *Neuroscience* 2007;150:522-36.
17. Buysse D, Reynolds C, Monk T, Berman S, Kupfer D. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193-213.
18. Agargun M, Kara H, Anlar O. The validity and reliability of Pittsburgh Sleep Quality Index. *Turkish Journal of Psychiatry* 1996;7:107-15.
19. Stewart AL, Hays RD, Ware JE. The MOS short-form general health survey. Reliability and validity in a patient population. *Med Care* 1988;26:724-35.
20. Kocyigit H. Reliability and validity of the Turkish version of short form-36 (SF-36) : a study in a group of patients with rheumatic diseases. *Turk J Drugs Ther* 1999;12:102-6.
21. Leja M, Grinberga-Derica I, Bilgicler C, Steininger C. Review: Epidemiology of *Helicobacter pylori* infection. *Helicobacter* 2019;24(Suppl 1):e12635. doi: 10.1111/HEL.12635
22. Liu Q, Zeng X, Wang W, Huang RL, Huang YJ, Liu S, Huang YH, Wang YX, Fang QH, He G, Zeng Y. Awareness of risk factors and warning symptoms and attitude towards gastric cancer screening among the

- general public in China: a cross-sectional study. *BMJ Open* 2019;9. doi: 10.1136/bmjopen-2019-029638
23. Cheng YH, Tung TH, Chen PE, Tsai CY. Risk of incident gastroesophageal reflux disease (GERD) in patients with sleep disorders: a population-based cohort study. *Sleep and Biological Rhythms* 2021;19:5-11.
  24. Mogavero MP, DelRosso LM, Fanfulla F, Bruni O, Ferri R. Sleep disorders and cancer: State of the art and future perspectives. *Sleep Med Rev* 2021;56. doi: 10.1016/j.smrv.2020.101409
  25. Orr WC, Fass R, Sundaram SS, Scheimann AO. The effect of sleep on gastrointestinal functioning in common digestive diseases. *Lancet Gastroenterol Hepatol* 2020;5:616-24.
  26. Voigt RM, Forsyth CB, Keshavarzian A. Circadian rhythms: a regulator of gastrointestinal health and dysfunction. *Expert Rev Gastroenterol Hepatol* 2019;13:411-24.
  27. Feyisa ZT, Woldeamanuel BT. Prevalence and associated risk factors of gastritis among patients visiting Saint Paul Hospital Millennium Medical College, Addis Ababa, Ethiopia. *PLoS One* 2021;16. doi: 10.1371/journal.pone.0246619
  28. Yang Q, Wang H, Shen H. A Case-control Study on Risk Factors for Chronic Gastritis in Yuhang District, Hangzhou. *Chinese Journal of Prevention and Control of Chronic Non communicable Diseases* 2006;14:27.
  29. Caruso CC, Lusk SL, Gillespie BW. Relationship of work schedules to gastrointestinal diagnoses, symptoms, and medication use in auto factory workers. *Am J Ind Med* 2004;46:586-98.
  30. Lee S, Chae CH, Park C, Lee HJ, Son J. Relationship of shift work with endoscopic gastritis among workers of an electronics company. *Scand J Work Environ Health* 2020;46:161-7.
  31. Guo S, Gao Q, Jiao Q, Hao W, Gao X, Cao JM. Gastric mucosal damage in water immersion stress: Mechanism and prevention with GHRP-6. *World J Gastroenterol* 2012;18:3145-55.
  32. Jiang H, Ling Z, Zhang Y, Mao H, Ma Z, Yin Y, Wang W, Tang W, Tan Z, Shi J, Li L, Ruan B. Altered fecal microbiota composition in patients with major depressive disorder. *Brain Behav Immun* 2015;48:186-94.
  33. Gebru D. Factors associated with gastric disease among students of Hawassa University: The case of collage of Agriculture students. *American Journal of Theoretical and Applied Statistics* 2018;7:207-14.
  34. Novaes Lipp ME, Lopes TM, Novaes Lipp LM, Falsetti MZ. Stress in Brazil. *International Journal of Psychiatry Research* 2020;3:1-4.
  35. Cho Y, Jung T, Jin H, Kim J, Kim D, Kim S. Relationship between Depression and Endoscopic Erosive Gastritis in Men and Women Aged over 40 Years. *Korean Journal of Family Practice* 2017;7:681-7.
  36. Cheng Y, Keita M, Si XB. Investigation on Risk Factors of Chronic Gastritis in the Population of Conakry in Guinea. *Indian Journal of Pharmaceutical Sciences* 2021;83:1-5.
  37. Firdous J, Muhamad N, Ab Latif N, Syazwani D, Hidayah N. A Descriptive Study on Lifestyle Factors Influencing Gastritis among University Students of UniKL RCMP in Malaysia. *International Bimonthly Indian Journal Of Natural Sciences [Internet]* (2016). [www.tnsroindia.org](http://www.tnsroindia.org). © IJONS, 6. [www.tnsroindia.org](http://www.tnsroindia.org).
  38. Filik L, Ozer N. Short sleep duration of overweight and obese patients with erosive esophagitis and gastritis. *Indian J Gastroenterol* 2015;34:408-9.